arXiv:nlin/0007004v1 [nlin.PS] 3 Jul 2000

Spatiotemporally localized solitons in resonantly absorbing Bragg reflectors

M. Blaauboer,[a,b] G. Kurizki,[a] and B.A. Malomed[b]
[a] *Chemical Physics Department, Weizmann Institute of Science,*
Rehovot 76100, Israel
[b] *Department of Interdisciplinary Studies, Faculty of Engineering,*
Tel Aviv University, Tel Aviv 69978, Israel

(February 3, 2008)

We predict the existence of spatiotemporal solitons ("light bullets") in two-dimensional self-induced transparency media embedded in a Bragg grating. The "bullets" are found in an approximate analytical form, their stability being confirmed by direct simulations. These findings suggest new possibilities for signal transmission control and self-trapping of light.

PACS numbers: 42.65.Tg, 78.66.-w, 42.65.Sf

Light propagation in periodic dielectric structures exhibits a variety of unique regimes that are technologically promising: nonlinear filtering, switching and distributed feedback amplification [1]. Of particular interest are gap solitons (GSs), i.e., moving or standing self-localized field structures centered in a band gap of the grating. These self-localized field structures arise due to the interplay between the medium nonlinearity and resonant Bragg reflections. Their spectrum is tuned away from the Bragg resonance by the nonlinearity at sufficiently high field intensities. Theoretical studies of gap solitons in Bragg gratings with Kerr nonlinearity [2] have been followed up by their experimental observation in a nonlinear optical fiber with the grating written on it [3]. Gap solitons have also been theoretically studied in gratings with second harmonic generation [4].

A principally different mechanism giving rise to gap solitons has been found in studies of models consisting of a periodic array of thin layers of *resonant two-level systems* (TLS's) separated by half-wavelength nonabsorbing dielectric layers, i.e., a *resonantly absorbing Bragg reflector* (RABR) [5–7] (Fig. 1). Such a RABR has been shown, for *any* Bragg reflectivity, to produce a vast family of stable solitons, both standing and moving ones. As opposed to the 2π solitons arising in self-induced transparency (SIT), i.e., near-resonant field-TLS interaction in a *uniform medium* [8–10], gap solitons in a RABR can have an *arbitrary* pulse area [5–7]. The existence of GS solutions can only be consistently demonstrated in a RABR with *thin* active TLS layers. By contrast, a recent attempt [11] to obtain such solutions in a periodic structure *uniformly* filled with active TLS's is physically doubtful, and fails for many parameter values.

The potential applications of GSs are based on the system's ability to filter out (by Bragg reflection) all pulses except for those satisfying the GS dispersion condition, as well as the control of pulse shape and velocity [5–7]. It would be clearly desirable to supplement these advantages by immunity to transverse diffraction of the pulse, i.e. to achieve *simultaneous* transverse as well as longitudinal self-localization in the structure. This calls for the consideration of "light bullets" (LBs), multi-dimensional solitons that are localized in both space and time. In the last decade they have been theoretically investigated in various nonlinear optical media [12–18], and the first experimental observation of a quasi two-dimensional (2D) bullet was recently reported [19]. In a recent work [20], we have predicted that uniformly 2D and 3D self-induced transparency (SIT) media can support stable light bullets.

In the present work, we aim to extend this investigation to RABR's of the kind shown in Fig. 1, so as to combine LB and GS properties in resonantly absorbing media. We find that a RABR with *any* Bragg reflectivity and *any* absorption cross-section can support the propagation of attenuated stable LBs, which are closely related to the bullets we have found in uniform SIT media [20]. It should be noted that 2D LBs supported by a combination of the Bragg reflector with a different (second-harmonic-generation) nonlinearity were theoretically investigated in Ref. [14].

We start by considering a 2D SIT medium with a spatially-varying refractive index $n(z,x)$, which is described by the lossless Maxwell-Bloch equations [9],

$$-i\mathcal{E}_{xx}+n^2\,\mathcal{E}_\tau+\mathcal{E}_z+i\,(1-n^2)\,\mathcal{E}-\mathcal{P}=0, \tag{1a}$$

$$\mathcal{P}_\tau-\mathcal{E}W=0, \tag{1b}$$

$$W_\tau+\frac{1}{2}(\mathcal{E}^*\mathcal{P}+\mathcal{P}^*\mathcal{E})=0. \tag{1c}$$

Here $\mathcal{E}$ and $\mathcal{P}$ are the slowly varying amplitudes of the electric field and polarization, W is the inversion, z and x are the longitudinal and transverse coordinates (measured in units of the resonant-absorption length), and τ is time (measured in units of the input pulse duration). The Fresnel number, which governs the transverse diffraction in the 2D and 3D propagation, has been incorporated in x and the detuning of the carrier frequency ω_0 from the central atomic-resonance frequency was absorbed in $\mathcal{E}$ and $\mathcal{P}$. The Fresnel number F, detuning $\Delta\Omega$, and wave vector $k_0\equiv\omega_0/c$ can be brought back into Eqs. (1) by the transformation $\mathcal{E}(\tau,z,x)\rightarrow(2/k_0)\;\mathcal{E}(\tau,z,x)\exp(-i\Delta\Omega\cdot\tau)$, $\mathcal{P}(\tau,z,x)\rightarrow$

$\left(4/k_0^2\right)\,\mathcal{P}(\tau,z,x)\exp(-i\Delta\Omega\tau)$, $W(\tau,z,x)\rightarrow\left(4/k_0^2\right)\,W$, $\tau\rightarrow(2/k_0)\,\tau$, $z\rightarrow(2/k_0)\,z$, and $x\rightarrow\sqrt{2/k_0F}\,x$. To neglect the polarization dephasing and inversion decay we assume pulse durations that are short on the time scale of the relaxation processes. Equations (1) are then compatible with the local constraint $|\mathcal{P}|^2+W^2=1$, which corresponds to the Bloch-vector conservation [9]. In the absence of the x-dependence and for $n(z,x)=1$, Eq. (1a) reduces to the sine-Gordon (SG) equation which has the soliton solution $\mathcal{E}(\tau,z)=2\alpha\,\mathrm{sech}(\alpha\tau-z/\alpha+\Theta_0)$, where α and Θ_0 are real parameters.



FIG. 1. Periodic modulation (2) of the refractive index in a structure of periodically alternating layers. Adapted from Ref. [5].

We proceed to search for LBs in a 2D medium subject to a resonant periodic longitudinal modulation, i.e., the above-mentioned RABR (Fig. 1). Accordingly, we assume the following periodic modulation of the refractive index [21]

$$n^2(z)=n_0^2\,\left[1+a_1\cos(2k_cz)\right]. \tag{2}$$

Here, n_0 and a_1 are constants and $k_c=\omega_c/c$, with ω_c being the central frequency of the band gap. A RABR is then constructed by placing very thin layers (much thinner than $1/k_c$) of two-level atoms, whose resonance frequency is close to ω_c, at the maxima of this modulated refractive index. We aim to consider the propagation of an electromagnetic wave with a frequency close to ω_c through a 2D RABR. Due to the Bragg reflections, the electric field $\mathcal{E}$ is decomposed into forward- and backward-propagating components $\mathcal{E}_F$ and $\mathcal{E}_B$, which satisfy equations that are a straightforward generalization of the 1D version in Refs. [5–7],

$$\begin{aligned}&-i\Sigma^+_{\tau xx}+i\Sigma^-_{zxx}+\Sigma^+_{\tau\tau}-\Sigma^+_{zz}\\&+\eta\Sigma^+_{xx}+\eta^2\Sigma^+-2\mathcal{P}_\tau-2i\eta\mathcal{P}=0, &&\text{(3a)}\\&-i\Sigma^-_{\tau xx}+i\Sigma^+_{zxx}+\Sigma^-_{\tau\tau}-\Sigma^-_{zz}\\&\quad-\eta\Sigma^-_{xx}+\eta^2\Sigma^-+2\mathcal{P}_z=0, &&\text{(3b)}\\&\quad\mathcal{P}_\tau+i\Delta\Omega\mathcal{P}-\Sigma^+W=0, &&\text{(3c)}\\&\quad W_\tau+\frac{1}{2}(\Sigma^{+*}\mathcal{P}+\Sigma^+\mathcal{P}^*)=0. &&\text{(3d)}\end{aligned}$$

Here $\Sigma^\pm\equiv 2\tau_0\mu n_0(\mathcal{E}_F\pm\mathcal{E}_B)/\hbar$, $\tau_0\equiv n_0\mu^{-1}\sqrt{\hbar/2\pi\omega_c\rho_0}$, μ is the transition dipole moment, and ρ_0 is the density of the two-level atoms. The parameter η is a ratio of resonant-absorption length in the two-level medium to the Bragg reflection length, and can be expressed as $\eta=a_1\omega_c\tau_0/4$.

In the 1D case, a family of *exact* soliton solutions to Eqs. (3) was found in Ref. [6]: $\Sigma^{(\pm)}=2\left(\alpha;-\sqrt{\alpha^2+2}\right)\mathrm{sech}\Theta(\tau,z)\cdot\exp\left(i\Phi\right)$, $\mathcal{P}=2\,\mathrm{sech}\Theta(\tau,z)\,\tanh\Theta(\tau,z)\cdot\exp\left(i\Phi\right)$, $W=2\mathrm{sech}^2\Theta(\tau,z)-1$, where $\Theta(\tau,z)\equiv\alpha\tau+\sqrt{\alpha^2+2}\,z+\Theta_0$, $\Phi\equiv\eta M\tau+\eta Nz+\phi$, $M\equiv-(\alpha^2+1)$, $N\equiv-\alpha\sqrt{\alpha^2+2}$, and $\Delta\Omega=\eta(\alpha^2+1)$. The shape of the fields Σ^+ and Σ^- in this solution is similar to the SG soliton in the uniform 1D SIT medium. Inspired by this analogy and the fact that there exist LBs in the uniform 2D SIT medium which reduces to the SG soliton in 1D [20], we search for a LB solution to the 2D equations (3), which reduces to the exact soliton in 1D. To this end, we try the following approximation,

$$\Sigma^+=2\,\alpha\,\sqrt{\mathrm{sech}\Theta_1\mathrm{sech}\Theta_2}\;e^{i\eta M\tau+i\eta Nz+i\pi/4}, \tag{4a}$$

$$\Sigma^-=-2\,\sqrt{\alpha^2+2}\,\sqrt{\mathrm{sech}\Theta_1\mathrm{sech}\Theta_2}\;e^{i\eta M\tau+i\eta Nz+i\pi/4}, \tag{4b}$$

$$\begin{aligned}\mathcal{P}=&\sqrt{\mathrm{sech}\Theta_1\mathrm{sech}\Theta_2}\{(\tanh\Theta_1+\tanh\Theta_2)^2+\\&\frac{1}{4}\alpha^2C^4[(\tanh\Theta_1-\tanh\Theta_2)^2-2(\mathrm{sech}^2\Theta_1+\\&\mathrm{sech}^2\Theta_2)]^2\}^{1/2}\;e^{i\eta M\tau+i\eta Nz+i\nu},\;W=\left[1-|\mathcal{P}|^2\right]^{1/2},\end{aligned} \tag{4c}$$

with $\Theta_1(\tau,z)\equiv\alpha\tau+\sqrt{\alpha^2+2}\,z+\Theta_0+Cx$, $\Theta_2(\tau,z)\equiv\alpha\tau+\sqrt{\alpha^2+2}\,z+\Theta_0-Cx$, the phase ν and coefficient C being real constants.

The ansatz (4) satisfies Eqs. (3a) and (3b) exactly, while Eqs. (3d) are satisfied to order $|\alpha|C^2$, which requires that $|\alpha|C^2\ll 1$. The ansatz is relevant for *arbitrary* η, admitting both weak ($\eta\ll 1$) and strong ($\eta>1$) reflectivities of the Bragg grating, provided that the detuning remains small with respect to the gap frequency, or $\eta\ll\omega_c/(\alpha^2+1)$. Comparison with numerical simulations of Eqs. (3), using (4) as an initial configuration, tests this analytical approximation and shows that it is indeed fairly close to a numerically exact solution; in particular, the shape of the bullet remains within 98% of its originally presumed shape after having propagated a large distance, as is shown in Fig. 2.



FIG. 2. The forward-propagating electric field of the 2D "light bullet" in the Bragg reflector, $|\mathcal{E}_F|$, vs. time τ and transverse coordinate x, after having propagated the distance $z = 1000$. The parameters are $\alpha = 1$, $C = 0.1$ and $\Theta_0 = -1000$. The field is scaled by the constant $\hbar/4\tau_0\mu n_0$.

We stress that 2D or 3D LB solutions of the variable-separated form $\Sigma^+ \sim \Sigma^- \sim f(\tau, z) \cdot g(x)$ do *not* exist in RABR. Indeed, substituting this into Eqs. (3a) and (3b) yields only a trivial solution of the form $\Sigma^\pm \sim \exp(iA\tau + iBx)$, with constant A and B.

We briefly discuss experimental conditions under which LBs can be observed in RABRs. The incident pulse should have uniform transverse intensity within its diameter d. For the transverse diffraction to be strong enough, one needs $\alpha_{\text{eff}} d^2/\lambda_0 < 1$, where α_{eff} and λ_0 are the inverse resonant-absorption length and carrier wavelength, respectively [22]. For $\alpha_{\text{eff}} \sim 10^3$ m^{-1} and $\lambda_0 \sim 10^{-4}$ m, one thus requires a diameter $d < 10^{-4}$ m, which implies that the transverse medium size $L_x \sim$ a few μm.

In order to realize a RABR, thin layers of *rare-earth ions* [23] embedded in a semiconductor structure with a spatially-periodic RI [24] may be used. The two-level atoms in the layers should be rare-earth-ions with density of $10^{15} - 10^{16}$ cm^{-3}, whose resonant-absorption time and inverse length are respectively $\tau_0 \sim 10^{-13} - 10^{-12}$ s and $\alpha_{\text{eff}} \sim 10^4 - 10^5$ m^{-1}. The parameter η can vary from 0 to 100 and the detuning is $\sim 10^{12} - 10^{13}$ s^{-1}. In a RABR with transverse size 10 μm, LBs depicted in Fig. 2 are localized on the time and transverse-length scales $\sim 10^{-13}$ s and 1 μm. Cryogenic conditions strongly extend the dephasing time T_2 and thus the LB lifetime, well into the μsec range [23]. The construction of suitable structures constitutes a feasible experimental challenge.

To conclude, we have studied light bullets in SIT media embedded in a resonantly-absorbing Bragg reflector. Light bullets in a multi-dimensional Bragg reflector have the potential of serving as a novel type of optical filters, which stably transmit selected signal frequencies through their spectral gap and block others. They can also be used to both spatially and temporarily localize light in certain frequency bands.

M.B. acknowledges support through a fellowship from the Israeli Council for Higher Education. Support from ISF, Minerva and BSF is acknowledged by G.K.

---

[1] See the *Photonic Band-Gap Bibliography,* compiled by J.P. Dowling and H.O. Everitt, at http://hwilwww.rdec.redstone.army.mil/MICOM/wsd/ST/RES/PBG/pbgbib.html

[2] A.B. Aceves *et al.*, Opt. Lett. **17**, 1566 (1992); B.A. Malomed and R.S. Tasgal, Phys. Rev. E **49**, 5787 (1994); I.V. Barashenkov *et al.*, Phys. Rev. Lett. **80**, 5117 (1998); A. De Rossi *et al.*, *ibid.* **81**, 85 (1998); A.R. Champneys, *et al.*, *ibid.* **80**, 4169 (1998).

[3] B.J. Eggleton *et al.*, Phys. Rev. Lett. **76**, 1627 (1996); C.M. de Sterke *et al.*, J. Lightwave Technol. **15**, 1494 (1997).

[4] T. Peschel *et al.*, Phys. Rev. E **55**, 4730 (1997); C. Conti *et al.*, Phys. Rev. Lett. **78**, 2341 (1997); Phys. Rev. E **57**, R1251 (1998); H. He and P. Drummond, Phys. Rev. Lett. **78**, 4311 (1997); S. Trillo and P. Ferro, Opt. Lett. **20**, 438 (1995); P. Drummond *et al.*, Opt. Comm. **123**, 394 (1996); H. He *et al.*, Phys. Rev. E **59**, 6064 (1999).

[5] A. Kozhekin and G. Kurizki, Phys. Rev. Lett. **74**, 5020 (1995).

[6] A. Kozhekin, G. Kurizki and B.A. Malomed, Phys. Rev. Lett. **81**, 3647 (1998).

[7] T. Opatrny, B.A. Malomed and G. Kurizki, Phys. Rev. E **60**, 6137 (1999).

[8] S.L. McCall and E.L. Hahn, Phys. Rev. Lett. **18**, 408 (1967).

[9] A.C. Newell and J.V. Moloney, *Nonlinear Optics* (Addison-Wesley: Redwood City, CA, 1992).

[10] A.A. Maimistov, A.M. Basharov, S.O. Elyutin, and Yu. M. Sklyarov, Phys. Rep. **191**, 1 (1990).

[11] N. Aközbek and S. John, Phys. Rev. E **58**, 3876 (1998).

[12] Y. Silberberg, Opt. Lett. **15**, 1281 (1990).

[13] K. Hayata and M. Koshiba, Phys. Rev. Lett. **71**, 3275 (1993); B.A. Malomed, P.D. Drummond, H. He, A. Berntson, D. Anderson, and M. Lisak, Phys. Rev. E **56**, 4725 (1997); D. Mihalache, D. Mazilu, B.A. Malomed, and L. Torner, Opt. Comm. **152**, 365 (1998); D. Mihalache, D. Mazilu, J. Dörring , and L. Torner, Opt. Comm. **159**, 129 (1999); A. Desyatnikov, A.I. Maimistov, and B.A. Malomed, Phys. Rev. E, in press.

[14] H. He and P. Drummond, Phys. Rev. E **58**, 5025 (1998).

[15] K.V. Kheruntsyan and P.D. Drummond, Phys. Rev A **58**, 2488 (1998); *ibid.* R2676 (1998).

[16] D.J. Frantzeskakis, K. Hizanidis, B.A. Malomed, and C. Polymilis, Phys. Lett. A **248**, 203 (1998).

[17] G. Gottwald, R. Grimshaw, and B.A. Malomed, Phys. Lett. A **248**, 208 (1998).

[18] P.D. Drummond, Opt. Comm. **49**, 219 (1984).

[19] X. Liu, L.J. Qian, and F.W. Wise, Phys. Rev. Lett. **82**, 4631 (1999).

[20] M. Blaauboer, B.A. Malomed, and G. Kurizki, Phys. Rev. Lett. **84**, 1906 (2000).

[21] In reality, $n^2(z)$ is represented by an infinite Fourier series. For the propagation of waves whose wavevector is close to k_c, however, only the coefficient a_1 is essential (see Ref. [7]).

[22] R.E. Slusher in: *Progress in Optics,* vol. 12, ed. by E. Wolf (North-Holland, Amsterdam, 1974).

[23] C. Greiner et al., Phys. Rev. A **60**, R2657 (1999).

[24] G. Khitrova, H.M. Gibbs, F. Jahnke, M. Kira, and S.W. Koch, Rev. Mod. Phys. **71**, 1591 (1999).